

2D Transformations

- Represent **linear** transformations by matrices
- To transform a point, represented by a vector, multiply the vector by the appropriate matrix.

2D Transformations

- Represent **linear** transformations by matrices
- To transform a point, represented by a vector, multiply the vector by the appropriate matrix.
- Recall the definition of matrix times vector:

$$\begin{bmatrix} a_{11}x + a_{12}y \\ a_{21}x + a_{22}y \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- A linear function $f(x)$ satisfies (by definition):

$$f(ax + by) = af(x) + bf(y)$$

- Note that “ x ” can be an abstract entity (e.g. a vector)—as long as addition and multiplication by a scalar are defined.
- Algebra reveals that matrix multiplication satisfies the above condition

- In particular., if we define $f(\mathbf{x}) = M \cdot \mathbf{x}$, where M is a matrix and \mathbf{x} is a vector, then

$$\begin{aligned} f(a\mathbf{x} + b\mathbf{y}) &= M(a\mathbf{x} + b\mathbf{y}) \\ &= aM\mathbf{x} + bM\mathbf{y} \\ &= af(\mathbf{x}) + bf(\mathbf{y}) \end{aligned}$$

- Where the middle step can be verified using algebra (next slide)

Proof that matrix multiplication is linear

$$\begin{aligned}
 M(a\mathbf{x} + b\mathbf{y}) &= \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix} \\
 &= \begin{bmatrix} a_{11}ax_1 + a_{11}by_1 + a_{12}ax_2 + a_{12}by_2 \\ a_{21}ax_1 + a_{21}by_1 + a_{22}ax_2 + a_{22}by_2 \end{bmatrix} \\
 &= a \begin{bmatrix} a_{11}x_1 + a_{12}x_2 \\ a_{21}x_1 + a_{22}x_2 \end{bmatrix} + b \begin{bmatrix} a_{11}y_1 + a_{12}y_2 \\ a_{21}y_1 + a_{22}y_2 \end{bmatrix} \\
 &= aM\mathbf{x} + bM\mathbf{y}
 \end{aligned}$$


- Now consider the linear transformation of a point on a line segment connecting two points, \mathbf{x} and \mathbf{y} .
- Recall that in parametric form, that point is: $t\mathbf{x} + (1 - t)\mathbf{y}$
- The transformed point is: $f(t\mathbf{x} + (1 - t)\mathbf{y}) = tf(\mathbf{x}) + (1 - t)f(\mathbf{y})$
- Notice that is a point on the line segment from the point $f(\mathbf{x})$ to the point $f(\mathbf{y})$,
- This shows that a linear transformation maps line segments to line segments

2D Transformations of objects

- To transform line segments, transform endpoints
- To transform polygons, transform vertices

2D Transformations

- Scale (stretch) by a factor of k




$$M = \begin{vmatrix} k & 0 \\ 0 & k \end{vmatrix}$$

($k = 2$ in the example)

2D Transformations

- Scale by a factor of (S_x, S_y)




$$M = \begin{vmatrix} S_x & 0 \\ 0 & S_y \end{vmatrix}$$

(Above, $S_x = 1/2$, $S_y = 1$)

2D Transformations

- Rotate around origin by θ (Orthogonal)




$$M = \begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix}$$

(Above, $\theta=90^\circ$)

2D Transformations

- Flip over y axis (Orthogonal)





$$M = \begin{vmatrix} -1 & 0 \\ 0 & 1 \end{vmatrix}$$

Flip over x axis is ?

2D Transformations

- Shear along x axis




$$M = \begin{vmatrix} 1 & a \\ 0 & 1 \end{vmatrix}$$

Shear along y axis is ?

2D Transformations

- Translation $(\mathbf{P}_{\text{new}} = \mathbf{P} + \mathbf{T})$



$\mathbf{M} = ?$

Homogenous Coordinates

- Represent 2D points by 3D vectors
- $(x,y) \rightarrow (x,y,1)$
- Now a multitude of 3D points (x,y,W) represent the same 2D point, $(x/W, y/W, 1)$
- Represent 2D transforms with 3 by 3 matrices
- Can now do translations
- Homogenous coordinates have other uses/advantages (later)

2D Translation in H.C.

$$\mathbf{P}_{\text{new}} = \mathbf{P} + \mathbf{T}$$

$$(x', y') = (x, y) + (t_x, t_y)$$

$$\mathbf{M} = \left| \begin{array}{ccc} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{array} \right|$$